T . F . . .

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington. VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it do not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)
29-05-2001	Journal Article	March 15, 2001
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER	
The Importance of High Reso Geometry in Modeling South	5b. GRANT NUMBER	
deciment, in measuring season		5c. PROGRAM ELEMENT NUMBER
		PE601153N
6. AUTHOR(S)		5d. PROJECT NUMBER
E. Joseph Metzg	5e TASK NUMBER	
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S	8. PERFORMING ORGANIZATION REPORT NUMBER	
Naval Research Laboratory		
Oceanography Division		
Stennis Space Center, MS		
		JA/7320-00-1000
9. SPONSORING / MONITORING AGENCY	10. SPONSOR/MONITOR'S ACRONYM(S)	
Office of Naval Research		
800 N. Quincy St.		11. SPONSOR/MONITOR'S REPORT
Arlington, VA 22217-5660	NUMBER(S)	

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release, distribution is unlimited

13. SUPPLEMENTARY NOTES

Geophysical Research Letters, Vol. 28, No. 6, Pages 1059-1062

20010808 027

14. ABSTRACT

As resolution is increased from 1/2 degrees to 1/32 degrees in Pacific Ocean simulations using the NRL Layered Ocean Model, marked changes are found in the Kuroshio's mean pathway as it intrudes into the South China Sea (SCS) via the Luzon Strait. With increased horizontal resolution comes a more accurate representation of the coastline geometry associated with the Batan/Babuyan Islands within the strait, and a reduction in the modeled westward intrusion of the Kuroshio into the SCS. The 1/16 degrees model is extremely sensitive to two very small scale shoals (Calayan Bank and a shoal north of Calayan Island) that are resolvable at this grid spacing. The exclusion of these three model gridpoints significantly alters the mean Kuroshio pathway to resemble the pathway from the 1/8 degrees model. In addition, excluding all islands within the Luzon Strait in the 1/16 degrees model gives a deep intrusion mean pathway as found in the 1/2 degrees model.

IS. SUBJECT TERMS

Luzon Strait, Layered Ocean Model, shoals

16. SECURITY CLASS	SIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON E. Joseph Meztger
a. REPORT Unclassified	b.AB\$TRACT Unclassified	c.THIS PAGE Unclassified	SAR	2	19b. TELEPHONE NUMBER (include area code) (228) 688-4762

PUBLICATION OR PRESENTATION RELEASE REQUEST

Division 7300 Pub Key: 2531

HEREFERENGES AND ENGLO	SURES.	2 TYPE OF PUBLICATION OR PRE	SENTATION	SIADMINISTRATIVE INFORMATION:
) Abstract only, not publishe	
Ref: (a) NRL Instruction 56 (b) NRL Instruction 55		() Book () Book Chapter	Route Sheet No.
(-)	/	() Conference Proceedings ((refereed)) Conference Proceedings (not refereed)	Job Order No.
Encl: (1) Two copies of sul	bject	() Invited speaker () Multimedia report	Classification II C
paper (or abstract)) Journal article (not refereed)	Connect of the
		() Other, explain) Oral Presentation, not publis	approval obtained ves
A AUTEORS				
Title of Paper or Presentation	1300 5 000 70 W	the second secon	e servicial de transferica de la como dela como de la como dela como de la como dela como de la com	and the state of t
Title of Paper of Fresentation	1011			
The Importance of High	h Horizon	tal Resolution and Accurate	Coastline Geometry in	n Modeling South China Sea Inflow
Author(s) Name(s) (First, I				a rivotaning double dilling and amily in
	,, <u></u> ,			
E. Joseph Metzger, Cod	e 7323; H	arley E. Hurlburt, Code 732	3	
It is intended to offer th	ie nanar t	n the		
it is intellued to offer th	iis hahei t	o tile	(Name of Con	ference)
			·	,
		(Date, Place and C	lassification of Conference)	
and/or for publication in	Geophys	ical Research Letters		
81		(Name and Classification of Publicat	-	(Name Of Publisner)
		pertinent publication/presen	tation data will be ente	red in the publications data base, in
accordance with referen				
It is the opinion of the a	luthor that	the subject paper (is)	(is not X) classified	, in accordance with reference (b).
communicated to the La	ate any di	sciosure of trade secrets or s	suggestions of outside	individuals or concerns which have been
This subject namer thas	i liporatory (never <u> </u>	ted in an official NRI AR	() contain and militarily critical technology
		never_/ been incorporal	ted in an official NRL/R	eporty
E. Joseph Metzger, Co		incipal Author)	_ C. page	(Signature)
Name and	Louis (F.	merpar Author)		Cognitute)
	فالمتكافئة فالمتكاف كالمتاب المعاد			
CODE	· ·	SIGNATURE	DATE	COMMENTS
Author(s) J. Metzger	18.7	mal Auton	9/14/00	
7.1.2.0051-	v		7570	
		· •		
	-	· · · · · · · · · · · · · · · · · · ·		
	.			
			1 1	
Section Head Rhodes or Acting	Mu	lt (Wh	9/14/60	
Branch Head			1///	
Harding or Acting		11 Km).	9/14/60	
Division Head				Release of this paper is approved. To the best knowledge of this Birings the outline.
		1 W/	9/15/	2. To the best knowledge of this Division, the subject matter of this paper (has)
Payne or Acting	(W	0/0	111/10	(has never I been classified.
Security, Code 7030.1			9/20/00	Paper of abstract was released. 2. A copy is filed in this office \$\$C-057-00
Office of Counsel,	1/1/	71/14 Pall 1		2. A copy is filed in this office 33.
Code 1008.3	1/1/6	la you	17/22/2K	
ADOR/Director NCST	,			
	1 1	, ,) , , , , , , , , , , , , , , , , ,	-,,	
ublic Affairs (Unclassified/ Inlimited Only), Code 7030.4	X) 11 /	o Valina)	9/00/00	
	XXXXX	()	7/40/00	
ivision, Code	ļ			
•	i		1	
uthor, Code			1 ·	, .

HQ-NRL 5219/1 (Rev. 5-97) (e) (Back)

The Importance of High Horizontal Resolution and Accurate Coastline Geometry in Modeling South China Sea Inflow

E. Joseph Metzger and Harley E. Hurlburt Naval Research Laboratory, Stennis Space Center, Mississippi

Abstract. As resolution is increased from 1/2° to 1/32° in Pacific Ocean simulations using the NRL Layered Ocean Model, marked changes are found in the Kuroshio's mean pathway as it intrudes into the South China Sea (SCS) via the Luzon Strait. With increased horizontal resolution comes a more accurate representation of the coastline geometry associated with the Batan/Babuyan Islands within the strait, and a reduction in the modeled westward intrusion of the Kuroshio into the SCS. The 1/16° model is extremely sensitive to two very small scale shoals (Calayan Bank and a shoal north of Calayan Island) that are resolvable at this grid spacing. The exclusion of these three model gridpoints significantly alters the mean Kuroshio pathway to resemble the pathway from the $1/8^{\circ}$ model. In addition, excluding all islands within the Luzon Strait in the $1/16^{\circ}$ model gives a deep intrusion mean pathway as found in the $1/2^{\circ}$ model.

Introduction

An accurate representation of numerical ocean model coastlines can have a significant impact on the mean pathways of ocean currents. Two issues at the core of proper model boundary definition are horizontal grid spacing and the accuracy of digital topographic databases. A finer horizontal grid allows for smaller scale features to be resolved, but many digital databases have serious deficiencies in defining small islands and shoals. Therefore, it is prudent to rely on other topographic sources, including the use of navigational charts.

Experience with the Naval Research Laboratory (NRL) Layered Ocean Model (NLOM) has shown that even relatively small changes in the model boundaries can have a significant impact on mean current pathways. These include a study on the effect of Halmahera Island on the Indonesia throughflow [Morey et al., 1999], modifications to the coastlines in the Intra-Americas Seas affecting the Florida Strait transport [Hurlburt and Townsend, 1994; Hurlburt and Hogan, 2000] and minor model boundary changes affecting the direction of the Sea of Japan outflow through the

This paper is not subject to U.S. copyright. Published in 2001 by the American Geophysical Union.

Paper number 2000GL012396.

Tsugaru Strait [Hurlburt et al., 1996].

Metzger and Hurlburt [1996; 2001] noted the modeled westward intrusion of the Kuroshio into the South China Sea (SCS) via the Luzon Strait was less pronounced with higher horizontal resolution. This was attributed in part to a better representation of the Luzon Strait island chain. This study focuses on the importance of high horizontal resolution and accurate coastline representation in modeling this region, but the implications extend beyond it.

Model Description

The NRL Layered Ocean Model is a primitive equation layered formulation where the equations have been vertically integrated through each layer. The 6-layer, thermodynamic, finite depth model of the Pacific Ocean used in this study is a descendant of the model by *Hurlburt and Thompson* [1980] and *Wallcraft* [1991]. Specific details and the model equations can be found in *Metzger and Hurlburt* [2001].

The model domain covers the Pacific Ocean from 20°S to 62°N and 109.125°E to 77.21°W. but the focus here is on the Luzon Strait that connects the northern SCS with the Pacific Ocean. Experiments with two different horizontal resolutions are used: 1/8° (1/8° in latitude by 45/256° in longitude for each prognostic model variable) and 1/16° (1/16° in latitude by 45/512° in longitude). All experiments are forced by a monthly climatology formed from European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis 10 m winds that span 1979-93 [Gibson et al., 1997]. The winds were converted to stresses using a stability dependent drag coefficient and temperature dependent air density [Kara et al., 2000]. No heat flux forcing is used in these experiments.

A modified version of the 1/12° ETOP05 bottom topography [NOAA, 1986] was used in the model. The 200 m isobath generally defines the model land/sea boundary but there are some exceptions, e.g. the Taiwan Strait, and the inflow/outflow straits to the Sea of Japan. A systematic problem of accurate representation of small islands exists in ETOP05, including the Batan and Babuyan Islands that run north-south within the Luzon Strait. Thus, navigational charts from the Defense Mapping Agency were also used to determine the model land/sea boundary in this and other




Plate 1. The bottom topography (in meters) and model coastline of the Luzon Strait in the a) 1/8° and b) 1/16° NLOM. The contour interval is 500 m.

regions [Defense Mapping Agency, 1980; Defense Mapping Agency, 1995]. Plate 1 depicts the bottom topography and model coastlines for the 1/8° and 1/16° models. Both contain the same islands and shoals within the Luzon Strait, although more crudely on the 1/8° grid.

Results/Discussion

Metzger and Hurlburt [1996; 2001] noted decreased westward intrusion of the modeled Kuroshio into the SCS with increasing horizontal resolution. In particular, the Batan and Babuyan Islands within the Luzon Strait were better represented as the resolution increased and no Luzon Strait islands were present in the 1/2° NLOM grid. Decreased westward intrusion is supported by a finding in Hurlburt and Thompson [1982] who used a constant absolute vorticity trajec-

tory analysis to determine a length scale $L_I = (v_c/\beta)^{1/2}$ for westward bending of the Loop Current in the Gulf of Mexico after it passes through the Yucatan Channel and is no longer supported by a western boundary. This westward bending is due to planetary vorticity (β) advection, where v_c is the velocity at the core of the current. Leonardi et al. [2001] additionally accounted for the inflow angle of the current as it separated from the coast and define the westward bending space scale $b_I = L_I(1 + \cos\theta_I)$, where θ_I is the inflow angle measured from the positive x-axis. This means that higher velocity in higher resolution models tends to increase the length scale for westward bending and to reduce the westward bending on a given length scale $< b_I$. The westward bending is further reduced by a narrower strait at higher resolution, a result consistent with the modeling work of Li et al., [1996].

While the northeast-southwest monsoon cycle causes large seasonal variability in the SCS, the focus here is on mean current pathways. Figure 1 shows six-year mean sea surface height (SSH) from $1/8^{\circ}$ and $1/16^{\circ}$ NLOM simulations. The simulations are identical in all respects except for the eddy viscosity (100 m^2/s for $1/8^{\circ}$ and 30 $\mathrm{m}^{2}/\mathrm{s}$ for $1/16^{\circ})$ and the horizontal resolution and hence the coastline configuration (Plate 1). The Kuroshio intrusion in the 1/8° model is much more pronounced than in the 1/16° model. Annual mean dynamic height at 100 m presented by Qu [2000] indicates a flow field more consistent with the 1/16° model simulation. The core of the Kuroshio parallels the Luzon Strait and doesn't exhibit significant westward intrusion in this $1/2^{\circ}$ analysis. The observed flow [Qu, 2000] that enters the SCS emanates from the lee of the Babuyan Islands in the southern half of the Luzon Strait, again a feature consistent with the $1/16^{\circ}$ model.

This difference between the two model simulations is attributed to the coastline representation within the




Plate 2. The 6-yr mean upper layer speed and currents from 1/16° NLOM simulations that a) include Calayan Bank and the shoal just north of Calayan Island and b) that exclude these three model gridpoints. The specified model gridpoints are highlighted in blue in panel a).






Figure 1. The 6-yr mean sea surface height from a) $1/8^{\circ}$ and b) $1/16^{\circ}$ NLOM simulations forced with climatological ECMWF 10 m winds. The contour interval is 2 cm. The net westward Luzon Strait transport is 2.2 Sv and 3.0 Sv for the $1/8^{\circ}$ and $1/16^{\circ}$ simulations, respectively. Qu [2000] estimates the Luzon Strait transport at 3.0 Sv.

Luzon Strait and the higher eddy viscosity at 1/8°. The coarser resolution 1/8° simulation cannot resolve the small scale islands and shoals as well as the 1/16° model. This is especially true in the southern half of the strait with regard to Calayan, Dalupiri and Fuga Islands and the passageways between them. This has a significant impact on the mean pathway of the Kuroshio's westward intrusion into the SCS (Figure 1). At 1/8° resolution, these shoals and islands are a contiguous group of land points. This effectively forces the core of the Kuroshio northward with the main branch entering the strait between Batan and Babuyan Islands, rather than between Babuyan and Camiguin Islands. At 1/16°, the resolution allows for an open passageway between Calavan and Dalupiri Islands and a lower eddy viscosity allows more flow through the passages south of Babuyan Island and the shoal north of Calayan Island. In this simulation the main branch of the Kuroshio takes a more southern route between Babuyan and Camiguin Islands.

To further isolate the cause of reduced Kuroshio intrusion into the SCS in the 1/16° model, note that two small scale shoals are having a substantial "blocking" effect. These are highlighted in blue on Plate 2a and are Calayan Bank (121.4°E, 19.7°N) and the shoal north of Calayan Island (121.5°E, 19.5°N). These shoals produce

two effects: 1) they narrow the Luzon Strait and thus reduce the westward bending compared to the case where they are absent, and 2) more importantly they deflect the flow so that the inflow angle is more northward, the latter corroborated by a 1/32° NLOM Pacific simulation (not shown). The significance of these three model gridpoints is illustrated in a simulation where the shoals were removed (Plate 2b). While the main branch of the Kuroshio still enters the strait between Babuyan and Camiguin Islands, the removal of the shoals at 1/16° changes the mean pathway to resemble that of the 1/8° model (Figure 1a). When Calayan Bank is removed in a 1/8° simulation, the mean pathway of the Kuroshio is almost unaffected (not shown) because the main branch of the Kuroshio continues to take the more northern route between Batan and Babuyan Islands and the inflow angle is essentially the same in both simulations.

What effect does the entire island chain have on Kuroshio intrusion? Figure 2 shows mean SSH from a 1/16° simulation with all Luzon Strait islands and shoals removed. Ocean depths at the surrounding gridpoints were averaged to convert the former model land points to ocean points. With a much wider Luzon Strait, the entire Kuroshio unrealistically loops into the northern SCS, a result similar to the 1/2° global model simulations of *Metzger and Hurlburt* [1996] that also excluded islands within the strait, and consistent with the idealized strait simulations by *Li et al.*, [1996].

Summary and Conclusions

Past and present simulations using the NRL Layered Ocean Model at $1/2^{\circ}$, $1/8^{\circ}$, $1/16^{\circ}$ and $1/32^{\circ}$ resolution have shown the importance of accurate representation of the north-south island chain within the Luzon Strait that connects the northern SCS and the Pacific Ocean. At $1/16^{\circ}$ and $1/32^{\circ}$ resolution, the mean pathway of the Kuroshio changes such that it bypasses the SCS rather than intruding deeply into it, a result consistent with the observations of Qu [2000]. The accuracy of




Figure 2. Same as Figure 1 except for a 1/16° NLOM simulation that excluded all Luzon Strait islands and shoals. The net westward Luzon Strait transport is 2.9 Sv.

model coastlines is obviously a function of horizontal resolution, but grid spacing as fine as $1/8^{\circ}$ cannot adequately resolve the Babuyan Islands in the southern latitudes of the Luzon Strait. At this resolution, the Calayan/Dalupiri/Fuga Island complex is a contiguous land mass and the higher eddy viscosity restricts the flow through open passages south of Babuyan Island. This forces the main branch of the Kuroshio to enter the SCS by a more northern route between Batan and Babuyan Islands and to enter with a substantial westward inflow component. In the 1/16° model, a passageway exists between Calayan and Dalupiri Island. This and the lower eddy viscosity allow the main branch of the Kuroshio to take a more southern route between Babuyan and Camiguin Islands. However, Calayan Bank has a "blocking effect" on this southern branch that turns the flow to a more northward inflow angle, thus reducing the westward bending into the SCS.

This has important consequences for modelers of this region. The results suggest that horizontal resolution of at least 1/16° is needed to adequately represent the islands and shoals in the Luzon Strait and thus simulate a Kuroshio more consistent with observations. Furthermore, the mean pathway is extremely sensitive to three seemingly inauspicious model gridpoints. Removal of these increases the westward intrusion to resemble 1/8° model results.

The need for accurate coastline representation goes beyond this region and resolution as demonstrated in earlier work cited in the introduction. Past experience with the NRL Layered Ocean Model has shown that interpolation of a standard digital topography like ETOP05 is insufficient since existing digital topographic databases are often not accurate at scales representable on grids at least as coarse as 1/4°. Thus, extra effort has been required when defining the land/sea grid, including the use of navigational charts and other topographic sources.

Acknowledgments. This work is part of the 6.1 Dynamics of Low Latitude Western Boundary Currents project sponsored by the Office of Naval Research under program element 601153N. The numerical simulations were performed on Cray T3Es at the Naval Oceanographic Office, Stennis Space Center, Miss. and the Engineer Research and Development Center, Vicksburg, Miss. using computer time from the Defense Dept. High Performance Computing Modernization Office. The NRL contribution number is JA/7320-00-1000.

References

Defense Mapping Agency, Balingtang Channel, Def.Map. Agency Hydrographic/Topographic Center Map 91ACO91160, 1980.

Defense Mapping Agency, Babuyan Channel, Def. Map. Agency Hydrographic/Topographic Center Map 91ACO91140, 1995.

Gibson, J.K., P. Kållberg, S. Uppala, A. Hernandez, A. Nomura, and E. Serrano, ECMWF Re-Analysis Project Report Series: 1. ERA description, ECMWF, Reading, Berkshire, UK, 72 pp, 1997.

Hurlburt, H.E. and P.J. Hogan, Impact of 1/8° to 1/64° resolution on Gulf Stream model-data comparisons in basinscale subtropical Atlantic Ocean models, Dyn. Atmos.

Ocean., 32, 283-329, 2000.

Hurlburt, H.E. and J.D. Thompson, A numerical study of Loop Current intrusions and eddy shedding, J. Phys. Oceanogr., 10, 1611-1651, 1980.

Hurlburt, H.E. and J.D. Thompson, The dynamics of the Loop Current and shed eddies in a numerical model of the Gulf of Mexico, In: Hydrodynamics of semi-enclosed seas, J.C. Nihoul (Ed.), pp. 243-298, Elsevier Scientific Publishing Company, Amsterdam, 1982.

Hurlburt, H.E., and T.L. Townsend, In: NRL effort in the North Atlantic, Data Assimilation and Model Evaluation Experiment - North Atlantic basin: Preliminary experiment plan, R.C. Willems, (Ed.), Tech. Rep. 2-95, pp. 30-35, Center for Ocean and Atmospheric Modeling, University of Southern Mississippi, Stennis Space Center, 1994.

Hurlburt, H.E., A.J. Wallcraft, W.J. Schmitz Jr., P.J. Hogan, and E.J. Metzger, Dynamics of the Kuroshio/Oyashio current system using eddy-resolving models of the North Pacific Ocean, J. Geophys. Res., 101, 941-976, 1996.

Kara, A.B., P.A. Rochford, and H.E. Hurlburt, Efficient and accurate bulk parameterizations of air-sea fluxes for use in general circulation models, J. Atmos. Ocean. Tech., 17, 1421-1438, 2000.

Leonardi, A.P, H.E. Hurlburt, E.J. Metzger and J.J. O'Brien, Dynamics of the North Hawaiian Ridge Current, J. Phys. Oceanogr., (submitted), 2001.

Li, W., Q. Liu, and S.-P. Cheng, The effect of a break in western boundary on the western boundary current, Acta Oceanographica Taiwanica, 35, 141-153, 1996.

Metzger, E.J. and H.E. Hurlburt, Coupled dynamics of the South China Sea, the Sulu Sea and the Pacific Ocean, J. Geophys. Res., 101, 12331-12352, 1996.

Metzger, E.J. and H.E. Hurlburt, The nondeterministic nature of Kuroshio penetration and eddy shedding in the South China Sea, J. Phys. Oceanogr., in press, 2001.

Morey, S.L., J.F.Shriver, and J.J. O'Brien, The effects of Halmahera on the Indonesian throughflow, J. Geophys. Res., 104, 23281-23296, 1999.

National Oceanic and Atmospheric Administration, ETOP05 digital relief of the surface of the Earth, Data Announce. 86-MGG-07, Natl. Geophys. Data Cent., Boulder, Colo, 1986.

Qu, T., Upper-layer circulation in the South China Sea, J. Phys. Oceanogr., 30, 1450-1460, 2000.

Wallcraft, A.J., The Navy Layered Ocean Model users guide, NOARL Rep. 35, 21 pp., Nav. Res. Lab., Stennis Space Center, Miss, 1991.

(Received Septémber 26, 2000; accepted January 3, 2001.)

E.J. Metzger and H.E. Hurlburt, Naval Research Laboratory, Code 7323, Stennis Space Center, MS 39529-5004. (email: metzger@nrlssc.navv.mil; hurlburt@nrlssc.navv.mil)